

David Fieleke, MD 601A NW O'Brien Road Lee's Summit, MO 64063 (816) 287-1528

Wound Care Instructions

• If you had a **biopsy**:

- 1. Keep the biopsy site dry the day of the procedure.
- 2. You may shower normally the next day, but do not immerse the wound in bath water.
- 3. Gently cleanse the wound 1-2 times daily with mild soap and water.
- 4. Pat dry, then apply a thick layer of Vaseline and cover with a Band-Aid.
- 5. Keep the wound moist with Vaseline to avoid scabbing.
- 6. It is normal for the wound to develop a thin yellowish film on its surface.
- 7. Continue wound care until the site fully heals; this may be 1-2 weeks depending on body location.

• If you were treated with **liquid nitrogen (freezing)**:

- 1. Blistering is a normal response. If blistering becomes bothersome, sterilize a needle over an open flame and pierce the blister to drain fluid out. Do not remove the blister as it protects the underlying skin.
- 2. Cleanse 1-2 times daily with mild soap and water.
- 3. Vaseline may be used for irritation or itching; it may also help speed healing time.
- 4. Continue wound care until site fully heals; this may be 1-2 weeks depending on body location.



David Fieleke, MD 601A NW O'Brien Road Lee's Summit, MO 64063 (816) 287-1528

Wound Care Instructions

• If you had a **biopsy**:

- 1. Keep the biopsy site dry the day of the procedure.
- 2. You may shower normally the next day, but do not immerse the wound in bath water.
- 3. Gently cleanse the wound 1-2 times daily with mild soap and water.
- 4. Pat dry, then apply a thick layer of Vaseline and cover with a Band-Aid.
- 5. Keep the wound moist with Vaseline to avoid scabbing.
- 6. It is normal for the wound to develop a thin yellowish film on its surface.
- 7. Continue wound care until the site fully heals; this may be 1-2 weeks depending on body location.

• If you were treated with **liquid nitrogen (freezing)**:

- 1. Blistering is a normal response. If blistering becomes bothersome, sterilize a needle over an open flame and pierce the blister to drain fluid out. Do not remove the blister as it protects the underlying skin.
- 2. Cleanse 1-2 times daily with mild soap and water.
- 3. Vaseline may be used for irritation or itching; it may also help speed healing time.
- 4. Continue wound care until site fully heals; this may be 1-2 weeks depending on body location.